

# Computational Experiments for the Problem of Hamiltonian Path with Fixed Number of Color Repetitions

**Anna Gorbenko**

Department of Intelligent Systems and Robotics  
Ural Federal University  
620083 Ekaterinburg, Russia  
gorbenko.ann@gmail.com

**Vladimir Popov**

Department of Intelligent Systems and Robotics  
Ural Federal University  
620083 Ekaterinburg, Russia  
Vladimir.Popov@usu.ru

## Abstract

In this paper we consider an approach to solve the problem of Hamiltonian path with fixed number of color repetitions for arc-colored digraphs. Our approach is based on usage of local search algorithms to solve a logical model for the problem.

**PACS:** 07.05.Fb, 07.05.Dz

**Keywords:** Hamiltonian path, arc-colored digraph, satisfiability problem, NP-complete

Let  $G = (V; E)$  be a digraph where

$$V = \{v[1], v[2], \dots, v[n]\}$$

is the set of nodes,

$$E = \{(v[i[1]], v[j[1]]), \dots\}$$

is the set of arcs. We assume that each arc of  $G$  has a color. If the number of colors of arcs is restricted by an integer  $c$ , we speak about  $c$ -arc-colored

digraphs. Let  $C[(u, v)]$  be the color of  $(u, v)$ . We can assume that

$$C[(u, v)] \in \{1, 2, \dots, c\}.$$

Let  $r$  be a fixed positive integer. Now, we consider the following problem.

THE PROBLEM OF HAMILTONIAN PATH WITH FIXED NUMBER OF COLOR REPETITIONS FOR  $c$ -ARC-COLORED DIGRAPHS:

INSTANCE: A  $c$ -arc-colored digraph  $G = (V; E)$ .

QUESTION: Is  $G$  has a Hamiltonian path

$$v[k[0]], v[k[1]], \dots, v[k[n-1]]$$

such that

$$\begin{aligned} C[(v[k[i]], v[k[i+1]])] = \\ C[(v[k[i+1]], v[k[i+2]])] = \dots = C[(v[k[i+r-1]], v[k[i+r]])], \\ |C[(v[k[j]], v[k[j+1]])] - C[(v[k[j+1]], v[k[j+2]])]| > 0, \end{aligned}$$

for all

$$\begin{aligned} i &= 0 \bmod r, \\ j+1 &= 0 \bmod r, \\ i &\in \{0, 1, \dots, n-1\}, \\ j &\in \{0, 1, \dots, n-1\}? \end{aligned}$$

Note that different Hamiltonian path problems are extensively studied (see e.g. [1] – [3]). have natural applications in the design of experiments. In particular, the problem of Hamiltonian path with fixed number of color repetitions for  $c$ -arc-colored digraphs can be used for scenarios creation for self-learning of intelligent mobile robots.

Note that the problem of Hamiltonian path with fixed number of color repetitions for  $c$ -arc-colored digraphs is **NP**-complete. Encoding different hard problems as instances of SAT and solving them with efficient satisfiability algorithms has caused considerable interest (see e.g. [4] – [19]). In this paper, we consider an approach to solve the problem of Hamiltonian path with fixed number of color repetitions for  $c$ -arc-colored digraphs. Our approach is based on an explicit reduction from the problem to the satisfiability problem. We consider different local search algorithms to solve a logical model for the problem.

Let

$$\begin{aligned} \varphi[1] &= \bigwedge_{0 \leq i \leq n-1} \bigvee_{1 \leq j \leq n} x[i, j], \\ \varphi[2] &= \bigwedge_{0 \leq i \leq n-1}, \quad (\neg x[i, j[1]] \vee \neg x[i, j[2]]), \\ &\quad 1 \leq j[1] < j[2] \leq n \end{aligned}$$

$$\begin{aligned}
\varphi[3] &= \bigwedge_{\substack{0 \leq i[1] < i[2] \leq n-1, \\ 1 \leq j \leq n}} (\neg x[i[1], j] \vee \neg x[i[2], j]), \\
\varphi[4] &= \bigwedge_{\substack{0 \leq i \leq n-2, \\ 1 \leq j[1] \leq n, \\ 1 \leq j[2] \leq n, \\ (j[1], j[2]) \notin E}} (\neg x[i, j[1]] \vee \neg x[i+1, j[2]]), \\
\varphi[5] &= \bigwedge_{\substack{0 \leq i \leq n-3, \\ 1 \leq j[1] \leq n, \\ 1 \leq j[2] \leq n, \\ 1 \leq j[3] \leq n, \\ C[(j[1], j[2])] \neq C[(j[2], j[3])], \\ i+1 \neq 0 \bmod r}} (\neg x[i, j[1]] \vee \neg x[i+1, j[2]] \vee \neg x[i+2, j[3]]), \\
\varphi[6] &= \bigwedge_{\substack{r-1 \leq i \leq n-r-1, \\ 1 \leq j[1] \leq n, \\ 1 \leq j[2] \leq n, \\ 1 \leq j[3] \leq n, \\ C[(j[1], j[2])] = C[(j[2], j[3])], \\ i+1 = 0 \bmod r}} (\neg x[i, j[1]] \vee \neg x[i+1, j[2]] \vee \neg x[i+2, j[3]]), \\
\xi &= \bigwedge_{i=1}^6 \varphi[i].
\end{aligned}$$

**Theorem.** *The function  $\xi$  is satisfiable if and only if there is a Hamilton path*

$$v[k[0]], v[k[1]], \dots, v[k[n-1]]$$

*in  $G$  such that*

$$\begin{aligned}
C[(v[k[i]], v[k[i+1]])] &= \\
C[(v[k[i+1]], v[k[i+2]])] &= \dots = C[(v[k[i+r-1]], v[k[i+r]])], \\
|C[(v[k[j]], v[k[j+1]])] - C[(v[k[j+1]], v[k[j+2]])]| &> 0,
\end{aligned}$$

*for all*

$$\begin{aligned}
i &= 0 \bmod r, \\
j+1 &= 0 \bmod r, \\
i &\in \{0, 1, \dots, n-1\}, \\
j &\in \{0, 1, \dots, n-1\}.
\end{aligned}$$

**Proof.** Let  $\xi = 1$ . In view of the definition of  $\xi$ , in this case, it is clear that  $\varphi[i] = 1$ , for all  $1 \leq i \leq 6$ .

Since  $\varphi[1] = 1$ , it is easy to see that for all  $0 \leq i \leq n-1$ , there is  $1 \leq j \leq n$  such that  $x[i, j] = 1$ . In view of  $\varphi[2] = 1$ , it is clear that for any  $0 \leq i \leq n-1$ , there is only one value of  $j$  such that  $x[i, j] = 1$ . Let  $x[i, a[i]] = 1$ . Since  $\varphi[3] = 1$ , it is easy to see that  $a[i] \neq a[j]$ , for any  $i \neq j$ . Therefore, values of

$x[i, j]$  can be considered as a choice of nodes for a path  $y[0], y[1], \dots, y[n-1]$ . In particular, we can assume that  $v[j]$  is the  $i$ th node in the path if and only if  $x[i, j] = 1$ .

Clearly,  $\varphi[4] = 1$  if and only if the sequence

$$y[0], y[1], \dots, y[n-1]$$

of nodes is a path. Since  $\varphi[5] = 1$ , it is easy to see that

$$C[(y[i], y[i+1])] = C[(y[i+1], y[i+2])] = \dots = C[(y[i+r-1], y[i+r])],$$

for all

$$i = 0 \bmod r,$$

$$i \in \{0, 1, \dots, n-1\}.$$

In view of  $\varphi[6] = 1$ , we obtain that

$$|C[(y[j], y[j+1])] - C[(y[j+1], y[j+2])]| > 0,$$

$$j+1 = 0 \bmod r,$$

$$j \in \{0, 1, \dots, n-1\}.$$

Now, we assume that there is a Hamilton path

$$v[k[0]], v[k[1]], \dots, v[k[n-1]]$$

in  $G$  such that

$$C[(v[k[i]], v[k[i+1]])] =$$

$$C[(v[k[i+1]], v[k[i+2]])] = \dots = C[(v[k[i+r-1]], v[k[i+r]])],$$

$$|C[(v[k[j]], v[k[j+1]])] - C[(v[k[j+1]], v[k[j+2]])]| > 0,$$

for all

$$i = 0 \bmod r,$$

$$j+1 = 0 \bmod r,$$

$$i \in \{0, 1, \dots, n-1\},$$

$$j \in \{0, 1, \dots, n-1\}.$$

Let  $x[i, j] = 1$  if and only if  $j = k[i]$ , for all  $0 \leq i \leq n-1$ . It is easy to check that  $\xi = 1$ . □

Using standard transformations (see e.g. [20]) we can easily obtain an explicit transformation  $\xi$  into  $\zeta$  such that  $\xi \Leftrightarrow \zeta$  and  $\zeta$  is a 3-CNF. It is clear that  $\zeta$  gives us an explicit reduction from the problem of Hamiltonian path with fixed number of color repetitions for  $c$ -arc-colored digraphs to 3SAT.

We have created a generator of natural instances for the problem of Hamiltonian path with fixed number of color repetitions for  $c$ -arc-colored digraphs. There is a well known site on which posted solvers for SAT [21]. We have used algorithms from [21]: fgrasp and posit. For solution of the problem, we have used a heterogeneous cluster. Each test was run on a cluster of at least 100 nodes. Note that due to restrictions on computation time (20 hours) we used savepoints. Selected experimental results are given in Table .

time	average	max	best
fgrasp	19.74 hr	56.04 hr	27.12 min
posit	12.63 hr	37.25 hr	31.08 min

Table 1: Experimental results for 3SAT

**ACKNOWLEDGEMENTS.** The work was partially supported by Analytical Departmental Program “Developing the scientific potential of high school” 8.1616.2011.

## References

- [1] A. Gorbenko and V. Popov, The Problem of Finding Two Edge-Disjoint Hamiltonian Cycles, *Applied Mathematical Sciences*, 6 (2012), 6563-6566.
- [2] A. Gorbenko and V. Popov, Hamiltonian Alternating Cycles with Fixed Number of Color Appearances, *Applied Mathematical Sciences*, 6 (2012), 6729-6731.
- [3] A. Gorbenko and V. Popov, The Hamiltonian Alternating Path Problem, *IAENG International Journal of Applied Mathematics*, 42 (2012) 204-213.
- [4] A. Gorbenko and V. Popov, Footstep Planning for Humanoid Robots, *Applied Mathematical Sciences*, 6 (2012), 6567-6571.
- [5] A. Gorbenko, M. Mornev, V. Popov, and A. Sheka, The problem of sensor placement for triangulation-based localisation, *International Journal of Automation and Control*, 5 (2011) 245-253.
- [6] A. Gorbenko and V. Popov, SAT Solvers for the Problem of Sensor Placement, *Advanced Studies in Theoretical Physics*, 6 (2012) 1235-1238.
- [7] A. Gorbenko and V. Popov, Clustering Algorithm in Mobile Ad Hoc Networks, *Advanced Studies in Theoretical Physics*, 6 (2012) 1239-1242.

- [8] A. Gorbenko, M. Mornev, and V. Popov, Planning a Typical Working Day for Indoor Service Robots, *IAENG International Journal of Computer Science*, 38 (2011) 176-182.
- [9] A. Gorbenko and V. Popov, The Problem of Selection of a Minimal Set of Visual Landmarks, *Applied Mathematical Sciences*, 6 (2012) 4729-4732.
- [10] A. Gorbenko and V. Popov, On the Problem of Placement of Visual Landmarks, *Applied Mathematical Sciences*, 6 (2012) 689-696.
- [11] A. Gorbenko and V. Popov, On the Longest Common Subsequence Problem, *Applied Mathematical Sciences*, 6 (2012), 5781-5787.
- [12] A. Gorbenko and V. Popov, On the Problem of Sensor Placement, *Advanced Studies in Theoretical Physics*, 6 (2012) 1117-1120.
- [13] A. Gorbenko and V. Popov, The Longest Common Parameterized Subsequence Problem, *Applied Mathematical Sciences*, 6 (2012), 2851-2855.
- [14] A. Gorbenko and V. Popov, The set of parameterized k-covers problem, *Theoretical Computer Science*, 423 (2012), 19-24.
- [15] A. Gorbenko and V. Popov, The Far From Most String Problem, *Applied Mathematical Sciences*, 6 (2012), 6719-6724.
- [16] A. Gorbenko and V. Popov, Multi-agent Path Planning, *Applied Mathematical Sciences*, 6 (2012), 6733-6737.
- [17] A. Gorbenko and V. Popov, Programming for Modular Reconfigurable Robots, *Programming and Computer Software*, 38 (2012) 13-23.
- [18] A. Gorbenko, V. Popov, and A. Sheka, Localization on Discrete Grid Graphs, *Lecture Notes in Electrical Engineering*, 107 (2012), 971-978.
- [19] A. Gorbenko and V. Popov, Task-resource Scheduling Problem, *International Journal of Automation and Computing*, 9 (2012), 429-441.
- [20] A. Gorbenko and V. Popov, The c-Fragment Longest Arc-Preserving Common Subsequence Problem, *IAENG International Journal of Computer Science*, 39 (2012), 231-238.
- [21] SATLIB — The Satisfiability Library. [Online]. Available: <http://people.cs.ubc.ca/~hoos/SATLIB/index-ubc.html>

**Received: December 3, 2012**